

Nkx2-5 mediates differential cardiac differentiation through interaction with Hoxa10

Ann N. Behrens¹, Michelina Iacovino^{1,3}, Jamie L. Lohr¹, Yi Ren¹, Claudia Zierold¹, Richard P. Harvey⁴, Michael Kyba^{1,3}, Daniel J. Garry^{1,2} and Cindy M. Martin^{1,2}.

¹Lillehei Heart Institute, University of Minnesota, Minneapolis, Minnesota

²Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, Texas.

³Department of Developmental Biology, University of Texas Southwestern Medical Center, Dallas, Texas

⁴Victor Chang Cardiac Research Institute, Lowy Packer Building, 405 Liverpool St., Darlinghurst 2010, Australia; and St. Vincent's Clinical School, University of New South Wales, Kensington 2052, Australia.

Running title: Hoxa10 inhibits cardiac differentiation

Keywords: Nkx2-5, Hoxa10, Cardiogenesis, Cardiac patterning, Cardiac progenitor cells, Mouse

Corresponding author:

Cindy M. Martin, MD
University of Minnesota
420 Delaware St. S.E. MMC 508, Minneapolis, MN 55455
Phone: (612) 625-6173
Fax: (214) 624-8118
E-mail: cmmartin@umn.edu

Abstract

The regulation of cardiac differentiation is complex and incompletely understood. Recent studies have documented that Nkx2-5 positive cells are not limited to the cardiac lineage but can give rise to endothelial and smooth muscle lineages. Other work has elucidated that, in addition to promoting cardiac development, Nkx2-5 plays a larger role in mesodermal patterning although the transcriptional networks that govern this developmental patterning are undefined. By profiling early Nkx2-5 positive progenitor cells, we discovered that the progenitor pools of the bisected cardiac crescent are differentiating asynchronously. This asymmetry requires Nkx2-5 as it is lost in the Nkx2-5 mutant. Surprisingly, the posterior Hox genes *Hoxa9* and *Hoxa10* were expressed on the right side of the cardiac crescent, independently of Nkx2-5. We describe a novel, transient and asymmetric cardiac-specific expression pattern of the posterior *Hox* genes, *Hoxa9* and *Hoxa10* and utilize the embryonic stem cell/embryoid body (ES/EB) model system to illustrate that *Hoxa10* impairs cardiac differentiation. We suggest a model whereby *Hoxa10* cooperates with Nkx2-5 to regulate the timing of cardiac mesoderm differentiation.

Keywords: Nkx2-5, Hoxa10, Cardiogenesis, Cardiac patterning, Cardiac progenitor cells, Mouse

Introduction

Congenital heart disease (CHD) affects approximately 1% of live births and accounts for the largest incidence of death due to birth defects in the first year of life [1,2]. Although numerous studies have defined the anatomical defects associated with CHD, the molecular networks that underpin these perturbations are incompletely defined [3,4]. Studies utilizing genetically modified mouse models have demonstrated the essential roles of transcription factors and signaling molecules at discrete stages of heart development [4-6]. *Nkx2-5* is one of the earliest markers of the cardiac lineage as it is abundantly expressed in the cardiac progenitor cells that form the cardiac crescent (E7.75) [7,8]. *Nkx2-5* is expressed throughout cardiac development and persists in the adult myocardium. Embryos lacking *Nkx2-5* are nonviable (E9.5-E10.5) due to growth retardation and gross abnormalities of the heart including a failure of ventricular chamber development [7,8]. Several recent studies have demonstrated that early *Nkx2-5* expressing progenitor cells are multipotent giving rise to cardiomyocyte, smooth muscle and endothelial lineages [9,10] and likewise, *Nkx2-5* knockout embryos have significant defects in these lineages [7,8,11]. These findings support the notion that *Nkx2-5* plays a larger role in mesodermal patterning and the early *Nkx2-5* expressing cells in the cardiac crescent may not yet be restricted exclusively to the cardiac lineage.

The Hox family of transcription factors are critical for early anterior-posterior (A-P) patterning of developing embryos and also regulate proliferation, differentiation and migration of multiple cell types [12,13]. Human and murine *HOX* genes are arranged in four clusters (A-D) and are positioned within each cluster in a 5'-3' fashion in the order in which they are expressed. The anterior groups, *Hox 1-8*, are homologous with the *Drosophila* antennapedia group (including *Abd-A*) while the *Abd-B* homologs consist of the posterior groups, *Hox 9-13* [14]. Studies in the hematopoietic system have identified that several *Hox* genes, including *Hoxa9* and *Hoxa10*, are expressed in the most primitive hematopoietic cells [15,16]. Enforced expression of *Hoxa10* has been shown to block differentiation and increase the number of early hematopoietic progenitors [16] and retroviral overexpression of *Hoxa10* not only dramatically impaired myeloid and lymphoid differentiation but resulted in the development of acute myeloid leukemia in a significant percentage of these mice [15]. The role

of *Hoxa9* and *Hoxa10* in the development of the reproductive track has been well described. *Hoxa10* is abundantly expressed in the urogenital system of both female and male mice [17]. Female mice with a targeted mutation of *Hoxa10* demonstrate urogenital deformities [18], and are sterile because of the absence of *Hoxa-10* expression in the adult uterus. *Hoxa10* mutant males display unilateral or bilateral cryptorchidism [17] which results in abnormal spermatogenesis and increasing sterility with age [17,19]. Although Hox genes typically were considered to be expressed only during embryonic development, the continued *Hoxa10* gene expression in the adult has been described in tissues with high plasticity such as bone marrow and endometrium [16,20]. In the reproductive tract the posterior Hox genes responsible for patterning the developing embryonic genitourinary tract are also responsible for remodeling the adult genitourinary tract as illustrated by the essential role of *Hoxa10* in the continuing developmental process of the female reproductive tract with each menstrual cycle and pregnancy [20]. Persistent Hox gene expression in the adult has been proposed as a mechanism by which the reproductive tract retains developmental plasticity [21].

The role of *Hox* genes in the cardiac lineage is less clear. Early studies in *Drosophila* noted a role for the *Hox* gene, *abd-A*, in cardiac tube development. The cardiac tube in flies consists of two regions, the anterior “aorta” and the posterior heart that contains contracting myocytes. In animals lacking *abd-A*, the heart myocytes failed to differentiate and mature appropriately, instead adopting an aorta-like phenotype [22,23]. Ectopic expression of *abd-A* however was sufficient to transform aortic myocytes into more posterior heart-like myocytes. Conversely, studies of *abd-B* (homolog of murine *Hox 9-13*) revealed that pan-mesodermal overexpression completely suppressed cardiogenesis, while *abd-B* loss-of-function mutants displayed formation of additional posterior myocardial cells [23,24]. Several anterior Hox genes have been implicated in mammalian cardiovascular development. A recent study identified severe cardiovascular defects in patients harboring a homozygous truncating mutation in *HOXA1* [25], however cardiovascular abnormalities were not identified in initial *Hoxa1* knockout mice models [26]. Given the newly described human mutation, more careful analysis revealed that almost three quarters of *Hoxa1* null mice had a cardiovascular defect [26]. Additional characterization of these mutant animals determined that *Hoxa1* is expressed in cardiac neural crest cells and is

required for neural crest specification [26,27]. *Hoxa3* expression has also recently been shown in cardiac neural crest cells and secondary heart field derived outflow tract myocardium; this expression is greatly enhanced with retinoic acid treatment [28]. The critical role of Hox genes in the A-P patterning of the developing embryo has been well described [29-32]. A spatial and temporal colinearity of Hox gene expression has been observed in the primitive streak and lateral mesoderm during the critical stages of early heart formation. Retinoic acid administration results in changes in patterning and expression of the Hox genes in chick limb buds and rat lung buds [33,34]. Similar RA-induced changes in A-P patterning of the primitive heart tube were observed in chick cardiogenic treated tissue [35,36]. These studies have led to the hypothesis that Hox genes are involved in early A-P patterning of the developing heart.

In the present study, we demonstrate that the *Nkx2-5* expressing progenitor cells of the cardiac crescent have an asymmetric gene expression profile. The asymmetry involves accelerated differentiation of left-sided progenitors (earlier expression of cardiomyocyte terminal markers on the left side), while the right side appears slower to differentiate maintaining higher expression of the early mesodermal marker *Brachyury* and the posterior *Hox* genes, *Hoxa9* and *Hoxa10*. Surprisingly, we find that the asymmetric expression of many transcripts requires functional *Nkx2-5*. We describe a novel, transient, early cardiac-specific expression pattern of the posterior *Hox* genes, *Hoxa9* and *Hoxa10* and utilize the embryonic stem cell/embryoid body (ES/EB) model system to illustrate that *Hoxa10* impairs cardiac differentiation. We also suggest a model whereby *Hoxa10* interacts with *Nkx2-5* to regulate the timing of cardiac mesoderm differentiation. Collectively, these studies enhance our understanding of molecular networks that govern discrete stages of early cardiac development.

Methods

FACS, transcriptome analyses

We utilized previously generated *Nkx2-5-promoter-EYFP* transgenic and *Nkx2-5* null mouse models [8,37] and isolated cardiac progenitors from the bisected E7.75 cardiac crescent following 0.25% Trypsin/EDTA (Invitrogen) digestion and FACS analysis [8,37]. Using a MoFlo Flow Cytometer (Beckman Coulter), EYFP labeled cells were collected directly into Tripure (Roche) and RNA was extracted and amplified as previously described [37]. Oligonucleotide array hybridizations were carried out according to the Affymetrix protocol as previously described [37,38]. cDNA synthesis and qRT-PCR were performed as previously described [38]. Expression was analyzed in triplicate by qRT-PCR using FAM-labeled Taqman probes from Applied Biosystems: *Gapdh* (Mn99999915_g1), *Tdglf1* (Mm01605855_g1), *Tnni* (Mm00441922_m1), *Kdr* (Mm00440099_m1), *Hoxa1* (Mm00439359_m1), *Hoxa2* (Mm00439361_m1), *Hoxa3* (Mm01326402_m1), *Hoxa4* (Mm01335255_m1), *Hoxa5* (Mm00439362_m1), *Hoxa6* (Mm00550244_m1), *Hoxa7* (Mm00657963_m1), *Hoxa9* (Mm00439364_m1), *Hoxa10* (Mm00433966_m1), *Hoxa11* (Mm00439360_m1), and *Hoxa13* (Mm00433967_m1) probes.

Analysis of Hoxa10 developmental expression pattern

Embryos were harvested and hearts isolated at designated developmental time points. In addition, at E8.25 (5-7 somite pairs), the heart tubes were isolated and the remaining embryos were divided into anterior segments minus heart and posterior segments. RNA extraction, cDNA synthesis and qRT-PCR reactions were performed as previously described [38]. Expression was analyzed in triplicate by qRT-PCR using FAM-labeled Taqman probes from Applied Biosystems: *Gapdh* (Mn99999915_g1), *Hoxa9* (Mm00439364_m1) and *Hoxa10* (Mm00433966_m1) probes.

Generation of an inducible ES/EB system for Hoxa10 overexpression

Doxycycline-inducible expression of *Hoxa10* in A2Lox mES cells was generated as previously described [39]. A2Lox cre mES cells lacking inserted *Hoxa10* plus doxycycline and *Hoxa10*-A2Lox mES cells without doxycycline were used as controls. The previously described reporter *Nkx2-5* emGFP ES cell line [40-42] was

infected with pSAM2-Hoxa10-mCherry or empty lentivirus along with pLenti-RtTA as previously described [43]. In brief, HA-*Hoxa10* cDNA was cloned into pSAM2-IRES-mCherry [43], a lentiviral construct with a doxycycline inducible promoter and the internal ribosome entry site (IRES) sequence followed by mCherry at the 3' end. Viral particles were prepared and cells were infected as previously described [43]. Infected cells were enriched by doxycycline induction followed by FACS for mCherry positive cells. During the last enrichment, cells were sorted by FACS for mCherry and SSEA1-APC (eBiosciences 51-8813-71) expression to enrich for virally integrated, undifferentiated ES cells. Expression was confirmed by flow cytometric analysis of mCherry, qRT-PCR using FAM-labeled Gapdh (Mn99999915_g1) and Hoxa10 (Mm00433966_m1) Taqman probes (Applied Biosystems) and Western blot using the following antibodies: goat-anti-Hoxa10 serum (1:200, Santa Cruz, Sc17158) and mouse-anti- α -tubulin serum (1:5000, Sigma T5168).

EB differentiation

ES cells were preplated for 30 minutes on gelatin coated flasks to remove the mouse embryonic fibroblasts (MEFs). Cells were resuspended at a concentration of 10,000 cells/ml in mouse EB differentiation medium (IMDM, 15% Fetal Bovine Serum, 2mM Glutamax, 1x Penicillin and Streptomycin, 450 μ M 1-thioglycerol (MTG), 200 μ g/ml Holo-Transferrin, and 50 μ g/ml Ascorbic acid). 10 μ l hanging drops were plated and cultured at 37°C with 5% CO₂ to induce differentiation. After 48 hours (at EB D2), hanging drops were washed from plates and further incubated in petri dishes on a rotating platform (70 rpm) at 37°C with 5% CO₂. Cells were fed with fresh medium with or without doxycycline every 48 hours, as previously described [44]. A2Lox cre mES cells lacking inserted Hoxa10 plus doxycycline and Hoxa10-A2Lox mES cells without doxycycline were used as controls. Gene expression was analyzed using qRT-PCR or Western blot analysis as previously described [39] using the following antibodies: goat-anti-Hoxa10 serum (1:200, Santa Cruz, Sc17158), rabbit-anti-cardiac Troponin I serum (1:3000, Abcam, ab47003), mouse-anti-cardiac Troponin T serum (1:3000, Abcam, ab8295), rabbit-anti-Conexin 43 serum (1:6000, Abcam, ab11370) and mouse-anti- α -tubulin serum (1:5000, Sigma, T5168). Day 4 EBs were dissociated, cells were incubated for 30 minutes on ice in the dark

with Flk-1-PE (eBiosciences 12-5821-83) and PDGFR α -APC (eBiosciences 17-1404-81). The Flk-1-PE, PDGFR α -APC double positive cardiac progenitors were quantified by FACS analysis on a BD FACS Aria [10, 45,46]. Nkx2-5-expressing cells were collected from day 4 EBs following 0.25% Trypsin/EDTA (Invitrogen) digestion and FACS analysis. qRT-PCR reaction was performed as previously described [38]. Expression was analyzed in triplicate by qRT-PCR using FAM-labeled Taqman probes from Applied Biosystems: Gapdh (Mm99999915_g1), Brachyury (Mm00436877_m1), Nkx2-5 (Mm00657783_m1), Gata4 (Mm00484689_m1), Isl1 (Mm00517585_m1), Tbx5 (Mm00803518_m1), Aldh1a2 (Mm00501306_m1), and Hes1 (Mm01342805_m1) probes.

Transcriptional Assays

Luciferase assays were performed as previously described [38]. Briefly, the 825bp *Nppa* promoter region containing Nkx2-5 and Gata4 binding sites was amplified by PCR and subcloned into the pGL3 vector to construct the *Nppa-luc* reporter using the following primers: For: 5'-GGACACGAGTCTTGGGAGGC-3' and Rev: 5'-GGGCACGATCTGATGTTTGC-3'. The 4219bp *Nkx2-5* promoter region containing Nkx2-5 and Gata4 binding sites was amplified by PCR and subcloned into the pGL3 vector to construct the *Nkx2-5-luc* reporter using the following primers: For: 5'- AGGTTCTCTTTGGCAGCAGGCATCTT-3' and Rev: 5'-GGGTTTCTTGGCTCAGGGTTTTGGAC -3'. COS7 cells were cultured in HyClone Dulbecco's modified Eagle's medium, High Glucose (ThermoFisher Scientific) supplemented with 10%. The cells were transfected with control (pGL3-Luc), pNkx2-5-Luc, or pNppa-Luc constructs with or without Nkx2-5, Gata4, and Hoxa10 overexpression plasmids ensuring equal amounts of total DNA [38]. Cells were harvested 24 hours after transfection and Luciferase activity was analyzed with Dual Luciferase System (Promega, Madison, WI) and normalized with the Renilla luciferase. Transfections were done in triplicate.

Statistical analysis

All p-values were calculated using Student's *t*-test analysis.

Results and Discussion

Molecular asymmetry in the cardiac crescent

We have previously employed an *Nkx2-5-EYFP* transgenic mouse model to identify the transcriptome of cardiac progenitor cells [37]. We identified transcripts that were enriched in progenitor cell populations throughout all stages. As expected, these included cardiac transcriptional regulators (*Nkx2.5*, *GATA4*, *HOP*, *Mef2C*, *myocardin*) and myocardial structural genes (*Myl4*, *Myl7*, *Tncc*, *Tnni1* and *Tnnt2*). We also noted that, compared to adult cardiomyocytes, the cardiac progenitor cell transcriptome consisted of an induction of the vascular, endocardial and signaling proteins, emphasizing the important role of these factors in fate determination, proliferation, patterning and cardiogenesis [37]. Several recent studies have expanded this finding, demonstrating that early *Nkx2-5* expressing progenitor cells are multipotent giving rise to not only cardiomyocytes, but also smooth muscle and endothelial lineages [9,10]. Utilizing this same *Nkx2-5-EYFP* transgenic mouse model and Affymetrix array technology, we examined the gene expression signature of cardiac progenitors that populated the left versus the right regions of the cardiac crescent of single embryos. We bisected the cardiac crescent (E7.75) and collected the EYFP positive cardiac progenitors from the left and right crescent of individual developing embryos using FACS analysis (Fig. 1A). Cell counts demonstrated that the number of EYFP-labeled cells collected from the left (mean = 2044, SD = 499, n = 5) and the right (mean 2263, SD 294, n = 5) sides of the crescent were comparable. Overall the molecular signatures of each side of the crescent were largely similar with only 70 transcripts being significantly differentially regulated in both of the tandem array results (Supplemental Table 1). 47 transcripts were enriched on the left side of the crescent versus 23 transcripts showing right-sided enrichment. Elegant studies have defined the role of *Nodal* and its interaction with *Cripto* to initiate the left-right asymmetric program by regulating downstream effectors [47]. In crescent stage embryos their expression is known to be restricted to the left side of the embryo [48] and accordingly, we observed left-sided enrichment of *Nodal*, *Cripto* and *Pitx2*. However, it was striking to note that the majority of the genes enriched on the left-sided cardiac crescent were myogenic genes such as *Myl4*, *Myh6*, *Tncc*, *Tnnt2* and *Tpm1*. There were also genes associated with differentiated endothelium (*Fus* and

VCAM) and smooth muscle (*Apeg1* and *Dia7*). These findings suggested that the left-sided program was enriched in transcripts associated with cardiovascular differentiation. The right side of the crescent, in contrast, was not enriched in genes for any specific tissue program but appeared to represent more undifferentiated mesoderm as illustrated by the enriched expression of *Brachyury*. These findings suggest that cardiovascular differentiation initiates asymmetrically on the left side of the crescent similar to the initiation of asymmetric genes expression on the left side of the node [49].

As discussed above, recent studies have described the role of anterior Hox genes in mammalian cardiovascular development [26-28]. Our array studies confirmed the expression of several anterior Hox genes in the early Nkx2-5 expressing cardiovascular progenitors (Supplemental Figure 1A). Unexpectedly, we detected the expression of posterior Hox genes within the cardiac crescent. The array findings also suggested a right-sided enrichment of the Hox genes within the cardiac crescent. qRT-PCR analysis of the Hox A family (family of Hox transcripts most frequently expressed in the array) confirmed the expression of the posterior Hox genes in EYFP positive cardiac progenitors from the left and right crescent of an additional embryo (Supplemental Figure 1B). *Hoxa9* and *Hoxa10* were enriched on the right side of the crescent both by array and qRT-PCR and this data suggested that these posterior Hox genes may play a role in cardiovascular development. In mammals, expression of *Hoxa9* and *Hoxa10*, first seen by *in situ* hybridization at the posterior limb bud stage, appeared to respect an anterior border posterior to the heart and were therefore previously not thought to be expressed in the developing heart [50]. The anterior expression border for *Hoxa9* had been described as 4-5 segments anterior to the hindlimb or at the last thoracic vertebrae [50]. The last lumbar vertebrae or first sacral vertebrae has been noted to be the most anterior expression level for *Hoxa10* [50]. We believe that this is the first description of a cardiac expression pattern for *Hoxa9* and *Hoxa10*. Although left-sided enrichment of *XHoxc-8* has been described in posterior lateral plate mesoderm in tail bud stage *Xenopus* embryos [51], there also has been no prior report of left-right asymmetric expression patterns of mammalian *Hox* genes.

Early asymmetric gene expression in the cardiac crescent is Nkx2.5 dependent

To further investigate the significance of this early asymmetric transcriptional program, we analyzed the same left-right transcriptional program in the *Nkx2-5* null cardiac crescent. We crossed the *Nkx2-5-EYFP* transgenic embryo into the *Nkx2-5* null background. As described previously, we bisected crescent stage embryos and utilized FACS to isolate the respective cell populations (Fig. 1A). No significant differences were observed in the number of EYFP positive cells recovered from the *Nkx2-5* null cardiac crescents compared to the wildtype crescents (Fig. 1A). The transcriptome of the *Nkx2-5* null crescent cells however was markedly different compared to the WT pattern (Fig. 1B). The array results were confirmed using qRT-PCR for selected transcripts (Fig. 1C). While the WT transcriptome was dominated by myogenic genes displaying left-sided enrichment, the *Nkx2-5* null crescent only displayed two transcripts with left-sided enrichment (Supplemental Table 2). While *Lefty2* maintained its expected asymmetric expression, the level of left-sided enrichment of *Pitx* was decreased when compared to WT. Even more striking was that the expression both *Nodal* and *Tdgfl* was largely equalized in the *Nkx2-5* null crescent. Previous studies have demonstrated that *Nodal* initiates and directly induces *Pitx2c* expression but *Nkx2-5* is important in the maintenance of *Pitx2* expression at later stages of development in the absence of *Nodal* signaling [52]. Our findings suggest that the role of *Nkx2-5* in maintaining the L-R asymmetric gene expression begins very early in cardiovascular development. Prior studies suggest a number of early patterning defects in the *Nkx2-5* null embryos - growth retardation, incomplete ventricular trabeculation, absence of the interventricular conduction ring, incomplete definition of the right and left ventricles and a failure to form endocardial cushions [7,8]. These anatomical perturbations were accompanied by decreased expression of *Myl2*, *Hand1* and *Nppa* in the *Nkx2-5* null ventricle compared to the WT control [7]. In our current study, *Nkx2-5* null crescents exhibited a loss of the left-sided enrichment for cardiac transcripts enriched on the left side of the WT crescent. *Nkx2-5* has been shown to transcriptionally activate a number of genes important in cardiac differentiation [53,54]. The result of the lack of this gene induction is highlighted in the *Nkx2-5* null embryos as they display defective development of the ventricles of the heart. The disruption of this program in the null crescent further supports the notion that *Nkx2-5* regulates the fate determination of future cardiovascular lineages in the earliest specified multi-potent cardiac progenitors. While the pattern of left-sided transcript enrichment was significantly altered in the null transcriptome, many

right-side enriched transcripts including *T*, *Fli* and the newly identified cardiac expression of *Hoxa9* and *Hoxa10* retained the patterns they exhibited in the WT crescent suggesting their regulation was independent of *Nkx2-5*. Given the apparent lack of differentiation on the right side of the crescent, the recently discovered importance of the anterior Hox genes, *Hoxa1* and *Hoxa3*, in cardiac development and the prior described role of *Hoxa10* in inhibiting hematopoietic differentiation we investigated whether the posterior Hox genes, *Hoxa9* and *Hoxa10* could be important in cardiovascular differentiation.

Hoxa9 and Hoxa10 are expressed in the early developing heart

Because *Hoxa9* and *Hoxa10* are broadly expressed throughout neuroectoderm and have previously been only noted to be expressed in mesoderm posterior to the thoracic vertebrae [50], we carefully confirmed this novel, putative anterior expression domain. The maintained asymmetrical expression of *Hoxa9* and *Hoxa10* in the *Nkx2-5* null crescent stage embryos suggests that the regulation of posterior *Hox* genes in mesodermal patterning is independent of *Nkx2-5*. Using qRT-PCR we found that *Hoxa9* and *Hoxa10* were not only expressed in the E7.75 EYFP-positive cardiac progenitors, but enriched when compared to the EYFP-negative cells harvested from the whole embryo (Fig. 2A). Division of E8.25 (5-7 somite pair) embryos into isolated heart tube, anterior embryo minus the heart, and posterior embryo also revealed continued enrichment of *Hoxa9* and *Hoxa10* in the heart (Fig. 2B). The cardiac progenitor specific expression of both *Hoxa9* and *Hoxa10* was noted to be transient as expression decreased rapidly as cardiac development proceeded (Fig. 2C). Expression of *Hoxa9* and *Hoxa10* in early GFP-positive, *Nkx2-5*-expressing cardiac progenitors isolated from crescent equivalent day 4 EBs generated from the *Nkx2-5* emGFP reporter ES cell line was confirmed on both the transcript and protein levels (Fig. 2 D & E).

Hoxa10 inhibits Cardiac Differentiation in Cell Culture and Embryoid Bodies

Previous studies have clearly demonstrated that *Hoxa10* plays an important role in regulating hematopoietic differentiation including the ability of *Hoxa10* to impair differentiation of mesodermally derived progenitor cells following specification[15,16]. No role for *Hoxa10* in cardiovascular mesodermal differentiation in a

mammalian system has yet been described. To further examine our hypothesis that *Hoxa10* played a role in early cardiovascular development, we complemented our *in vivo* studies utilizing the *Nkx2-5-EYFP* transgenic embryos with the generation of a doxycycline-inducible *Hoxa10* overexpressing embryonic stem (ES) cell line. Our strategy is schematized in Supplemental Figure 2 and highlights that the reverse tetracycline transactivator (rtTA) binds to the tetracycline responsive element (TRE) to induce *Hoxa10* expression only after the addition of doxycycline. The day 4 EB cardiac progenitor population was significantly reduced in the *Hoxa10* overexpressing EBs compared to control EBs (Fig. 3A; 15 +/- 9% vs. 44 +/- 8%, n=8, p <0.05). Assays also revealed a significant decrease in the percentage of beating embryoid bodies at EB day eight in the *Hoxa10* overexpressing EBs compared to control EBs (Fig. 3B; 13 +/- 10% vs. 78 +/- 22%, n=8, p <0.05). Protein harvested from day 12 EBs revealed an absence of cardiac troponin I in the *Hoxa10* overexpressing EBs compared to doxycycline treated control EBs (Fig. 3C). This control confirmed that the decrease of beating and lack troponin expression was not simply a doxycycline effect. Protein harvested from *Hoxa10* day 12 EBs without doxycycline confirmed that without induction of *Hoxa10*, EBs generated from the doxycycline-inducible *Hoxa10* cell line were able to differentiate into cardiomyocytes as evidenced by robust expression of cardiac troponin I and cardiac troponin T (Fig. 3D & E). However, induction of *Hoxa10* overexpression again resulted in an absence of cardiac troponin I and cardiac troponin T (Fig. 3D & E). Preserved expression of connexin 43 in the *Hoxa10* day 12 EB protein samples supported that the lack of beating was not due to a defect in excitation or contraction coupling but rather a perturbation of cardiomyocyte differentiation as evidenced by the lack of cardiac structural proteins. The alteration in cardiogenesis did not appear to be a general posterior Hox gene effect as *Hoxa13* had a much lower effect on the percentage of Flk-1 and PDGFR α double positive cardiac progenitor cells (30.5 +/- 1.4% vs. 40 +/- 6.7%, n=4, p < 0.05) and no significant reduction in beating EBs (77 +/-27% vs. 62 +/- 32%, n = 4, p = 0.5) compared to control EBs.

Our initial studies in the *Hoxa10*-overexpressing EBs suggest that early *Hoxa10* misexpression restricts the specification of cells to a cardiac lineage and also impairs the differentiation of early progenitor cells into differentiated, contractile cardiomyocytes. To further address the role of *Hoxa10* on the *Nkx2-5* expressing

early progenitors of the cardiac crescent, we utilized the Nkx2-5 emb-GFP-Hoxa10-mCherry pSAM2 ES cell line in which lentiviral mediated Hoxa10 expression is induced with the addition of doxycycline in ES cells harboring an Nkx2-5-emGFP reporter. Using these engineered ES cells, Hoxa10 expression was induced from EB day 3-4 to correlate with increased Hoxa10 expression at the E7.75 (cardiac crescent) developmental time point. The total percentage of cardiac progenitor cells was again significantly reduced in the Hoxa10 overexpressing cells (Fig. 3G; 49 +/- 7% vs. 36 +/- 7%, n=5, p <0.05). However, there was no significant difference in the total percentage of GFP-positive, Nkx2-5 expressing cells in the Hoxa10 overexpressing cells versus uninduced cells (24 +/- 8% vs. 30 +/-9%, n=5, p > 0.05). Further analysis of these GFP-positive, Nkx2-5 expressing cells revealed a significant change in the characteristics of this population. In the uninduced sample, 71 +/- 5% of the GFP-positive-Nkx2-5 expressing cells were double positive for Flk-1 and PDGFR α , thus mostly representing cardiac progenitor cells (Fig. 3H). However, when Hoxa10 was overexpressed, the percentage of Flk-1 and PDGFR α -double positive Nkx2-5-expressing cells was significantly reduced (49 +/- 10%, n = 5, p < 0.05; Fig. 3H). qRT-PCR analysis of these GFP-positive, Nkx2-5 expressing cells collected at EB day 4 confirmed increased *Brachyury* expression and revealed a reduction in expression of the cardiac transcription factors *Nkx2-5*, *Gata-4*, *Tbx-5*, and *Isl-1* as well as a small decrease in *Aldh1a2* with overexpression of *Hoxa10*; *Hes1* expression, however, was not significantly altered (Fig. 3I). Thus, early transient overexpression of Hoxa10 does not appear to reduce the percentage of initial Nkx2-5 cardiovascular progenitors. The overexpression of Hoxa10 does however, appear to impair the differentiation of these Nkx2-5 expressing cells to a more cardiac restricted phenotype as illustrated by their reduced percentage of Flk-1/PDGFR α -double positive cardiac progenitors from day 4 EBs. Increased *Brachyury* expression and reduced expression of the cardiac transcription factors from these GFP-positive Nkx2-5 expressing cells following Hoxa10 overexpression further supports this hypothesis. We theorize that these cells most likely parallel the less differentiated Hoxa10 expressing cells of the right side of the cardiac crescent.

Hoxa10 interacts with Nkx2-5

The final set of studies was focused on defining the mechanism through which *Hoxa10* perturbed early cardiogenesis. The Hox protein family has been previously shown to interact with homeodomain proteins including Pbx and Meis family members [55]. Hox genes have the ability to act as both transcriptional activators and repressors [56] and recent work has suggested that these effects may be mediated through sequestering other proteins [57]. Given that the expression of several cardiac transcription factors in the EB derived early cardiac progenitors was significantly decreased with *Hoxa10* overexpression, and *Nkx2-5* in combination with *Gata4* is known to cooperatively transactivate the promoters of several genes important in cardiogenesis [53,54,58], we questioned if *Hoxa10* was acting as a cardiac transcriptional repressor. Transcriptional assays confirmed that the addition of *Hoxa10* decreased the *Nkx2-5*-*Gata4* mediated activation of the both the *Nkx2-5* and *Nppa* promoter (Supplemental Fig. 3 and Fig. 3J). This finding suggested a mechanism in which *Hoxa10* may be more broadly regulating cardiogenesis as *Nkx2-5* and *Gata4* are known to function as mutual cofactors for a number of genes important in cardiac development [53,54,58]. This further supports that *Hoxa10* plays an important role in specification and differentiation of the early cardiac progenitors.

Collectively, these data support a model whereby cardiac differentiation occurs in a left to right temporal wave across the *Nkx2-5* expressing cells of the cardiac crescent. Continued expression of *Hoxa10* impedes cardiac differentiation of the *Nkx2-5* positive early cardiac progenitor cells in the EB system. This is the first study that examines the asymmetric molecular program of the cardiac crescent and is the first to describe the expression and functional role of *Hoxa10* during early cardiac development. Definition of these transcriptional networks in cardiac progenitor cells will enhance our understanding of mesodermal patterning and cardiac differentiation during normal and perturbed cardiac development.

Acknowledgements

We thank Zhaohui Xu for his assistance with the pSAM2 lentiviral system generation. We also thank Alessandro Magli, Teresa Gallardo, and Amanda Massino for critical discussions and technical assistance. This work was supported by the National Institutes of Health (K08 HL102157 and U01 HL100407) and the American Heart Association (Jon Holden DeHaan Foundation 0970499).

Disclosures

None

References

1. Hoffman JI. (1995). Incidence of congenital heart disease: II. Prenatal incidence. *Pediatr Cardiol* 16:155-165.
2. Olson EN and D Srivastava. (1996). Molecular pathways controlling heart development. *Science* 272:671-676.
3. Harvey RP. (2002). Patterning the vertebrate heart. *Nat Rev Genet* 3:544-556.
4. Olson EN and MD Schneider. (2003). Sizing up the heart: development redux in disease. *Genes Dev* 17:1937-1956.
5. Brand T. (2003). Heart development: molecular insights into cardiac specification and early morphogenesis. *Dev Biol* 258:1-19.
6. Evans SM, D Yelon, FL Conlon and ML Kirby. (2010). Myocardial lineage development. *Circ Res* 107:1428-1444.
7. Tanaka M, Z Chen, S Bartunkova, N Yamasaki and S Izumo. (1999). The cardiac homeobox gene *Csx/Nkx2.5* lies genetically upstream of multiple genes essential for heart development. *Development* 126:1269-1280.
8. Lyons I, LM Parsons, L Hartley, R Li, JE Andrews, L Robb and RP Harvey. (1995). Myogenic and morphogenetic defects in the heart tubes of murine embryos lacking the homeo box gene *Nkx2-5*. *Genes Dev* 9:1654-1666.

9. Christoforou N, RA Miller, CM Hill, CC Jie, AS McCallion and JD Gearhart. (2008). Mouse ES cell-derived cardiac precursor cells are multipotent and facilitate identification of novel cardiac genes. *J Clin Invest* 118:894-903.
10. Bondue A, S Tannler, G Chiapparo, S Chabab, M Ramialison, C Paulissen, B Beck, R Harvey and C Blanpain. (2011). Defining the earliest step of cardiovascular progenitor specification during embryonic stem cell differentiation. *J Cell Biol.* 192:751-765.
11. Caprioli A, N Koyano-Nakagawa, M Iacovino, X Shi, A Ferdous, RP Harvey, EN Olson, M Kyba and DJ Garry. (2011). Nkx2-5 represses Gata1 gene expression and modulates the cellular fate of cardiac progenitors during embryogenesis. *Circulation.* 123:1633-1641.
12. Adams MD, SE Celniker, RA Holt, CA Evans, JD Gocayne, PG Amanatides, SE Scherer, PW Li, RA Hoskins, RF Galle, RA George, SE Lewis, S Richards, et al. (2000). The genome sequence of *Drosophila melanogaster*. *Science* 287:2185-2195.
13. Miller DF, SL Holtzman, A Kalkbrenner and TC Kaufman. (2001). Homeotic Complex (Hox) gene regulation and homeosis in the mesoderm of the *Drosophila melanogaster* embryo: the roles of signal transduction and cell autonomous regulation. *Mech Dev* 102:17-32.
14. Shah CA, H Wang, L Bei, LC Plataniias and EA Eklund. (2011). HoxA10 regulates transcription of the gene encoding transforming growth factor beta2 (TGFbeta2) in myeloid cells. *J Biol Chem.* 286:3161-3176.
15. Bromleigh VC and LP Freedman. (2000). p21 is a transcriptional target of HOXA10 in differentiating myelomonocytic cells. *Genes Dev* 14:2581-2586.
16. Magnusson M, AC Brun, N Miyake, J Larsson, M Ehinger, JM Bjornsson, A Wutz, M Sigvardsson and S Kalsson. (2007). HOXA10 is a critical regulator for hematopoietic stem cells and erythroid/megakaryocyte development. *Blood* 109:3687-3696.
17. Satokata I, G Benson and R Maas. (1995). Sexually dimorphic sterility phenotypes in Hoxa10-deficient mice. *Nature* 374:460-463.
18. Benson GV, H Lim, BC Paria, I Satokata, SK Dey and RL Mass. (1996). Mechanisms of reduced fertility in Hoxa-10 mutant mice: uterine homeosis and loss of maternal Hoxa-10 expression. *Development* 122:2687-2696.

19. Rijli FM, R Matyas, M Pellegrini, A Dierich, P Gruss, P Dolle and P Chambon. (1995). Cryptorchidism and homeotic transformations of spinal nerves and vertebrae in Hoxa-10 mutant mice. *Proc Natl Acad Sci U S A* 92:8185-8189.
20. Zanatta A, AM Rocha, FM Carvalho, RM Pereira, HS Taylor, EL Motta, EC Barcat and PC Serfini. (2010). The role of the Hoxa10/HOXA10 gene in the etiology of endometriosis and its related infertility: a review. *J Assist Reprod Genet* 27:701-710.
21. Du H and HS Taylor. (2004). Molecular regulation of mullerian development by Hox genes. *Ann N Y Acad Sci* 1034:152-165.
22. Ponzielli R, M Astier, A Chartier, A Gallet, P Therond and M Semeriva. (2002). Heart tube patterning in *Drosophila* requires integration of axial and segmental information provided by the Bithorax Complex genes and hedgehog signaling. *Development* 129:4509-4521.
23. Lovato TL, TP Nguyen, MR Molina and RM Cripps. (2002). The Hox gene abdominal-A specifies heart cell fate in the *Drosophila* dorsal vessel. *Development* 129:5019-5027.
24. Kim YO, SJ Park, RS Balaban, M Nirenberg and Y Kim. (2004). A functional genomic screen for cardiogenic genes using RNA interference in developing *Drosophila* embryos. *Proc Natl Acad Sci U S A* 101:159-164.
25. Tischfield MA, TM Bosley, MA Salih, IA Alorainy, EC Sener, MJ Nester, DT Oystreck, WM Chan, C Andrews, RP Erickson and EC Engle. (2005). Homozygous HOXA1 mutations disrupt human brainstem, inner ear, cardiovascular and cognitive development. *Nat Genet* 37:1035-1037.
26. Makki N and MR Capecchi. (2012). Cardiovascular defects in a mouse model of HOXA1 syndrome. *Hum Mol Genet* 21:26-31.
27. Makki N and MR Capecchi. (2011). Identification of novel Hoxa1 downstream targets regulating hindbrain, neural crest and inner ear development. *Dev Biol* 357:295-304.
28. Diman NY, S Remacle, N Bertrand, JJ Picard, S Zaffran and R Razsohazy. (2011). A retinoic acid responsive Hoxa3 transgene expressed in embryonic pharyngeal endoderm, cardiac neural crest and a subdomain of the second heart field. *PLoS One*. 6:e27624.
29. Davis AP, DP Witte, HM Hsieh-Li, SS Potter and MR Capecchi. (1995). Absence of radius and ulna in mice lacking hoxa-11 and hoxd-11. *Nature* 375:791-795.

30. Kondo T, P Dolle, J Zakany and D Duoule. (1996). Function of posterior HoxD genes in the morphogenesis of the anal sphincter. *Development* 122:2651-2659.
31. Podlasek CA, D Duboule and W Bushman. (1997). Male accessory sex organ morphogenesis is altered by loss of function of Hoxd-13. *Dev Dyn* 208:454-465.
32. Roberts DJ, RL Johnson, AC Burke, CE Nelson, BA Morgan and C Tabin. (1995). Sonic hedgehog is an endodermal signal inducing Bmp-4 and Hox genes during induction and regionalization of the chick hindgut. *Development* 121:3163-3174.
33. Cardoso WV, SA Mitsialis, JS Brody and MC Williams. (1996). Retinoic acid alters the expression of pattern-related genes in the developing rat lung. *Dev Dyn* 207:47-59.
34. Tabin CJ. (1991). Retinoids, homeoboxes, and growth factors: toward molecular models for limb development. *Cell* 66:199-217.
35. Yutzey KE and D Bader. (1995). Diversification of cardiomyogenic cell lineages during early heart development. *Circ Res* 77:216-219.
36. Yutzey KE, JT Rhee and D Bader. (1994). Expression of the atrial-specific myosin heavy chain AMHC1 and the establishment of anteroposterior polarity in the developing chicken heart. *Development* 120:871-883.
37. Masino AM, TD Gallardo, CA Wilcox, EN Olson, RS Williams and DJ Garry. (2004). Transcriptional regulation of cardiac progenitor cell populations. *Circ Res* 95:389-397.
38. Martin CM, A Ferdous, T Gallardo, C Humphries, H Sadek, A Caprioli, JA Garcia, LI Szweda, MG Garry and DJ Garry. (2008). Hypoxia-inducible factor-2alpha transactivates Abcg2 and promotes cytoprotection in cardiac side population cells. *Circ Res* 102:1075-1081.
39. Iacovino M, C Hernandez, Z Xu, G Bajwa, M Prather and M Kyba. (2009). A conserved role for Hox paralog group 4 in regulation of hematopoietic progenitors. *Stem Cells Dev* 18:783-792.
40. Hsiao EC, Y Yoshinga, TD Nguyen, SL Musone, JE Kim, P Swinton, I Espeneda, C Manalac, PJ deJong and BR Conklin. (2008). Marking embryonic stem cells with a 2A self-cleaving peptide: a NKX2-5 emerald GFP BAC reporter. *PLoS One* 3:e2532.
41. Zhang L, A Magli, J Catanese, Z Xu, M Kyba and RC Perlingeiro. (2011). Modulation of TGF-beta signaling by endoglin in murine hemangioblast development and primitive hematopoiesis. *Blood* 118:88-97.

42. Bosnakovski D, Z Xu, EJ Gang, CL Galindo, M Liu, T Simsek, HR Garner, S Agha-Mohammadi, A Tassin, F Coppee, A Belayew, RR Perlingeiro and M Kyba. (2008). An isogenetic myoblast expression screen identifies DUX4-mediated FSHD-associated molecular pathologies. *EMBO J* 27:2766-2779.
43. Darabi R, RW Arpke, S Irion, JT Dimos, M Grskovic, M Kyba and RC Perlingeiro. (2012). Human ES- and iPS-derived myogenic progenitors restore DYSTROPHIN and improve contractility upon transplantation in dystrophic mice. *Cell Stem Cell* 10:610-619.
44. Koyano-Nakagawa N, J Kweon, M Iacovino, X Shi, TL Rasmussen, L Borges, LM Zirbes, T Li, RC Perlingeiro, M Kyba and DJ Garry. (2012). Etv2 is expressed in the yolk sac hematopoietic and endothelial progenitors and regulates Lmo2 gene expression. *Stem Cells* 30:1611-1623.
45. Lindsley RC, JG Gill, TL Murphy, EM Langer, M Cai, M Mashayekhi, W Wang, N Niwa, JM Nerbonne, M Kyba and KM Murphy. (2008). Mesp1 coordinately regulates cardiovascular fate restriction and epithelial-mesenchymal transition in differentiating ESCs. *Cell Stem Cell* 3:55-68.
46. Sakurai H, T Era, LM Jakt, M Okada, S Nakai and S Nishikawa. (2006). In vitro modeling of paraxial and lateral mesoderm differentiation reveals early reversibility. *Stem Cells* 24:575-586.
47. Norris, DP and EJ Robertson. (1999). Asymmetric and node-specific nodal expression patterns are controlled by two distinct cis-acting regulatory elements. *Genes Dev* 13:1575-1588.
48. Adachi H, Y Saijoh, K Mochida, S Ohishi, H Hashiguchi, A Hirao and H Hamada. (1999). Determination of left/right asymmetric expression of nodal by a left side-specific enhancer with sequence similarity to a lefty-2 enhancer. *Genes Dev* 13:1589-1600.
49. Tabin CJ. (2006). The key to left-right asymmetry. *Cell* 27:27-32.
50. Burke AC, CE Nelson, BA Morgan and C Tabin. (1995). Hox genes and the evolution of vertebrate axial morphology. *Development* 121:333-346.
51. Thickett C and R Morgan. (2002). Hoxc-8 expression shows left-right asymmetry in the posterior lateral plate mesoderm. *Gene Expr Patterns* 2:5-6.
52. Shiratori H, R Sakuma, M Watanabe, H Hashiguchi, K Mochia, Y Sakai, J Nishino, Y Saijoh, M Whitman and H Hamada. (2001). Two-step regulation of left-right asymmetric expression of Pitx2: initiation by nodal signaling and maintenance by Nkx2. *Mol Cell* 7:137-149.

53. Durocher D, F Charron, R Warren, RJ Schwartz and M Nemer. (1997). The cardiac transcription factors Nkx2-5 and GATA-4 are mutual cofactors. *EMBO J* 16:5687-5696.
54. Lee Y, T Shioi, H Kasahara, SM Jobe, RJ Wiese, BE Markham and S Izumo. (1998). The cardiac tissue-restricted homeobox protein Csx/Nkx2.5 physically associates with the zinc finger protein GATA4 and cooperatively activates atrial natriuretic factor gene expression. *Mol Cell Biol* 18:3120-3129.
55. Sarno JL, HJ Kliman and HS Taylor. (2005). HOXA10, Pbx2, and Meis1 protein expression in the human endometrium: formation of multimeric complexes on HOXA10 target genes. *J Clin Endocrinol Metab* 90:522-528.
56. Svingen T and KF Tonissen. (2006). Hox transcription factors and their elusive mammalian gene targets. *Heredity (Edinb)* 97:88-96.
57. Shah N and S Sukumar. (2010). The Hox genes and their roles in oncogenesis. *Nat Rev Cancer* 10:361-371.
58. Reecy JM, X Li, M Yamada, FJ DeMayo, CS Newman, RP Harvey and RJ Schwartz. (1999). Identification of upstream regulatory regions in the heart-expressed homeobox gene Nkx2-5. *Development* 126:839-849.

Figure Legends

Figure 1. Hoxa9 and Hoxa10 are expressed in early cardiac progenitor cells of the cardiac crescent.

(A) Individual WT and Nkx2-5 null cardiac crescent stage embryos were bisected and EYFP-positive cells were collected from the right and left side of the crescent. Left WT mean = 2044, SD = 499, n = 5; left null mean = 1967, SD = 200, n = 4; right WT mean = 2263, SD = 294, n = 5; right null mean = 2167, SD = 252, n = 4. (B) Transcriptome analysis of the WT and Nkx2-5 null left vs. right side of the crescent. Note a significant induction of cardiac structural transcripts in the left vs. right WT crescent. This left-sided enrichment is absent in the Nkx2-5 null crescent. The right-sided enrichment in expression is largely unaltered in the absence of Nkx2-5. Significant differential transcript expression is denoted by green highlight (up-regulated) or red highlight (down-regulated). (C) qRT-PCR of selected transcripts in WT (black bar) and null (open bar) right and left crescent.

Figure 2. Characterization of cardiac Hoxa10 expression.

(A) Analyses of enrichment of Hoxa9 and Hoxa10 in the EYFP-positive Nkx2-5 expressing cells vs. EYFP-negative cells collected from the cardiac crescent using qRT-PCR; *, $p < 0.05$, $n = 3$. (B) qRT-PCR reveals Hoxa9 and Hoxa10 are enriched in the E8.25 (5-7 somite) heart compared to the remaining segments of the embryo; *, $p < 0.05$, $n = 3$. (C) Cardiac Hoxa9 and Hoxa10 expression is rapidly downregulated during cardiac development as assessed by qRT-PCR; *, $p < 0.05$, $n = 3$. (D) Analyses of fold change in the expression of the Hoxa9, Hoxa10 and Nkx2-5 from GFP-positive Nkx2-5 expressing cells vs. GFP-negative obtained from EBs harvested at day 4 using qRT-PCR; *, $p < 0.05$, $n = 3$. (E) Western analysis of EB day 4 GFP-positive Nkx2-5 expressing cells confirm Hoxa10 protein expression.

Figure 3. Hoxa10 inhibits cardiogenesis.

(A) Percentage of EB day 4, Flk-1-PDGFR α double positive cardiac progenitor cells are significantly decreased following Hoxa10 overexpression for 48hrs (day 2-4) of EB formation; 15 +/- 9% vs. 43.6 +/- 7.9%, *, $p < 0.05$, $n = 8$. (B) Reduction of beating in EBs overexpressing Hoxa10 following induction with doxycycline (Dox) on days 3-6 vs. control EBs plus Dox; 12.5 +/- 10.3 vs. 78.4 +/- 21.7, * $p < 0.05$, $n = 8$. (C) Western analysis reveals absence of cTroponin I in day 12 EBs overexpressing Hoxa10 following the induction with doxycycline (Dox) on days 3-6 vs. day 12 control EBs treated with doxycycline (Dox) on days 3-6). Western analysis reveals absence of cTroponin I (D) and cTroponin T (E) in day 12 Hoxa10 doxycycline-inducible EBs following the induction with doxycycline (Dox) on days 3-6 vs. no doxycycline. (F) Protein expression of Connexin 43 is preserved in day 12 EBs following Hoxa10 induction. (G) Nkx2-5-GFP reporter ES cell line was infected with Hoxa10 or empty lentivirus. Overexpression of Hoxa10 for 24 hrs (day 3-4 of EB formation) in cardiac crescent equivalent progenitors results in a reduction of total Flk-1/PDGFR α double positive cardiac progenitor cells; 35.5 +/- 7% vs. 48.6 +/- 7%; *, $p < 0.05$, $n = 5$. (H) Percentage of Nkx2-5 expressing GFP positive cells double positive for Flk-1 and PDGFR α is also significantly decreased with Hoxa10 overexpression; 49 +/- 10% vs. 71 +/- 5%; *, $p < 0.05$, $n = 5$. (I) qRT-PCR analyses revealed increase in Brachyury expression and reduction in the cardiac transcription factors *Nkx2-5*, *Gata-4*, *Tbx-5*, and *Isl-1* as

well as a small decrease in *Aldh1a2* in GFP positive cells collected from day 4 Nkx2-5-GFP EBs following the induction of Hoxa10 with doxycycline (Dox) for 24hrs (day3-4); *, $p < 0.05$, $n = 3$. Hes1 expression, however, was not significantly altered. (J) The Nppa promoter was fused to the luciferase (luc) reporter gene and was transfected into Cos7 cells with and without addition of the Nkx2-5, Gata4 or Hoxa10 expression plasmids. Empty pGL3 and Nppa promoter driven luciferase expression is shown. Nppa promoter revealed a 10.65 +/- 0.44 fold activation with Nkx2-5 which was reduced to 7.41 +/- 0.50 with addition of Hoxa10 and a 20.55 +/- 0.49 fold activation with Nkx2-5 and Gata4 which was reduced to 12.62 +/- 0.17 with addition of Hoxa10; *, $p < 0.05$, $n = 3$.